

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

**Environmental and spatial factors drive diatom species distribution in  
Alpine streams: Implications for biomonitoring**

**This is the author's manuscript**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/1704544> since 2023-01-27T10:25:28Z

*Published version:*

DOI:10.1016/j.ecolind.2019.105441

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

1 **Environmental and spatial factors drive diatom species distribution in Alpine streams:**  
2 **implications for biomonitoring**

3 Elisa Falasco<sup>1</sup>, Francesca Bona<sup>1</sup>, Catia Monauni<sup>2</sup>, Agnese Zeni<sup>2</sup>, Elena Piano<sup>1\*</sup>

4 <sup>1</sup> DBIOS, University of Turin, Via Accademia Albertina 13, I-10123, Turin, Italy

5 <sup>2</sup>APPA – Settore tecnico per la tutela dell’ambiente – U.O. Acqua, Via Lidorno 1, I-38123, Trento,  
6 Italy

7 \*Corresponding author: elena.piano@unito.it

8

9 **ABSTRACT**

10 Given their documented capacity to track changes in environmental conditions and human alterations,  
11 benthic diatom communities are at present widely used in biomonitoring programs to evaluate stream  
12 water quality. However, it is becoming more and more evident that species ecological preferences are  
13 not the only drivers of diatom community composition, since dispersal-related processes also play a  
14 role. This is particularly compelling in Alpine streams, where orographic conformation and human-  
15 related impacts limit dispersal of organisms. In addition, several environmental variables may  
16 influence diatom community in pristine or impacted sites. We here investigate the differential role of  
17 environmental and spatial factors in driving the community assemblages of diatoms in streams of the  
18 Eastern Italian Alps, focusing on both taxonomic and functional composition. We analysed data from  
19 110 samples collected on two different geological substrates, i.e. calcareous and siliceous, during the  
20 last eight years of biomonitoring programs, among which 64 collected in reference sites and 46 in  
21 impacted sites. We first evaluated whether diatom communities in reference and impacted sites are  
22 differentially shaped by environmental and spatial factors, highlighting the major role of spatial  
23 constraints in both of them. In particular, anthropogenic disruption of longitudinal connectivity in  
24 streams likely shaped impacted communities, as demonstrated by the increasing abundance of motile

25 taxa, which are associated with physical disturbance. Conversely, reference communities were mostly  
26 affected by spatially structured environmental variables, especially those related to streambed  
27 lithology. We then compared the taxonomic and functional composition of diatom communities  
28 between the two geological substrates in both reference and impacted sites to better highlight the  
29 differential role of this factor. Our results demonstrate that lithology strongly drives diatom  
30 community composition in reference but not in impacted sites, confirming our previous observations.  
31 The analysis of functional traits, however, highlighted how differences were due not only to the  
32 geological substrates, but also to other environmental variables, like flow velocity. Overall, the effect  
33 of the spatial component on the structure of diatom assemblages can represent a background noise in  
34 the framework of the river quality assessment, and this should be taken into account especially in  
35 those countries, like Italy, covering a broad range of mountain areas.

36 **Key-words:** reference sites, ecological guilds, functional groups, dispersal limitation, variation  
37 partitioning

38

## 39 **Highlights**

- 40 • Diatoms track changes in environmental conditions and human alterations
- 41 • Dispersal is expected to affect diatom distribution in Alpine streams
- 42 • Substrate lithology affects diatom community in reference sites
- 43 • Chemical variables and spatial processes affect diatom community in impacted sites
- 44 • Biomonitoring programs should account for the combined effects of spatial and
- 45 environmental processes

## 1. INTRODUCTION

Given their documented capacity to track changes in environmental conditions and human alterations, diatoms have been gradually included in all the European monitoring programs as biological indicators for the quality assessment of lotic freshwater ecosystems, after the publication of the Waterframe Directive (2000/60/CE). Diatom responses to environmental variables acting at multiple scales of analysis have been widely documented in literature (Rimet et al., 2004; 2007; 2009; Tison et al., 2005; Grenier et al., 2006; Bona et al., 2007; Soininen, 2007). For instance, Soininen (2007) identified flow velocity and substratum composition as key determinants of diatom composition at local scale, whereas ion concentrations and trophic conditions mainly drive changes in river diatom communities at a regional scale (Bona et al., 2007; Soininen, 2007). However, it has been recently proved that species ecological preferences alone are not sufficient to explain diatom community composition, since dispersal processes also play an important role in shaping communities (Heino, 2013).

Owing to their physical, topographic and climatic heterogeneity, Alpine streams are extremely heterogeneous, encompassing large variation in environmental variables, like flow velocity, temperature and ion concentrations, over short distances (Ward, 1994). This environmental heterogeneity likely exerts a selective pressure on organisms inhabiting Alpine lotic freshwater ecosystems, which track their optimal environmental conditions according to their ecological niche. In Alpine streams, where ion and nutrient concentrations are usually very low, geology and, as a consequence, conductivity and hardness resulted among the most important factors in shaping diatom communities (Rimet et al., 2004; 2007; 2009; Tison et al., 2005; Grenier et al., 2006), but also altitude, dissolved oxygen, nitrate, calcium, turbidity and pH are among the main environmental factors shaping diatom assemblages in such oligotrophic ecosystems (see Falasco et al., 2012 for an example in Western Alps and Cantonati, 1998; Cantonati et al., 2001; 2006; 2007; 2009; Beltrami et al. 2012 for examples in the Eastern Alps). However, recent works have highlighted that the expected

71 responses of diatoms to environmental parameters are not completely fulfilled, especially in mountain  
72 streams, due to the effect of spatial processes (Bottin et al., 2014; Dong et al., 2016).

73 Although composition of diatom communities is clearly shaped by environmental variables and  
74 habitat heterogeneity at local scale (i.e. species sorting), the role of spatial factors should be kept into  
75 account at regional/global scale (Soininen, 2007; Verleyen et al., 2009). The importance of diatom  
76 dispersal-related processes in the framework of the biological monitoring has been recently proved  
77 in lacustrine ecosystems (Vilmi et al., 2016). Although lake communities are more spatially-  
78 structured than stream ones, due to the higher dispersal limitation characterizing the lentic species  
79 (Soininen, 2007; Marquardt et al., 2017), Bottin et al. (2014) remarked the importance of the spatial  
80 component also in mountain streams, where steep slopes surrounding streams might limit organism  
81 dispersion (Dong et al., 2016). In addition, also human alterations such as stream channelization, flow  
82 regulations and the construction of reservoirs, interrupt habitat connectivity and limit dispersal of  
83 organisms. For instance, water scarcity, caused by both local impacts and global changes, has recently  
84 become the most critical threat for diatom lotic assemblages in mountain streams, altering the  
85 taxonomical and functional composition of benthic communities ((Bona et al., 2008; Falasco et al.,  
86 2018; Piano et al., 2019). Thus, anthropogenic impacts in Alpine streams are expected to alter not  
87 only chemical variables but also the spatial processes underlying the composition of diatom  
88 communities, overriding the role of natural environmental filters or physical barriers (such as  
89 lithological substrates and orographic barriers). A better understanding of the role of environmental  
90 and dispersal-related processes in shaping diatom communities is then compelling to enhance the  
91 implementation of diatom-based biomonitoring programs in Alpine streams.

92 One of the main outcomes of anthropogenic disturbance on the composition of biotic communities is  
93 biotic homogenization, which is defined as an increase in the similarity of species composition (Olden  
94 et al., 2004). Environmental filters and dispersal barriers caused by the human activities likely cause  
95 the loss of rare and specialised species, and the gain of widespread tolerant ones, with consequent

96 biotic homogenization. In addition, according to the “habitat templet theory” (Southwood, 1977,  
97 1988; Townsend and Hildrew, 1994), the above mentioned selection processes may have major  
98 effects on particular functional traits, like morphological or physiological attributes, or they could  
99 affect the realized niche of a certain species (Webb et al., 2010), causing also functional  
100 homogenization (Olden et al., 2004). Thus, the investigation of these processes may provide  
101 information on the effects of anthropogenic pressures on biotic communities.

102 According to Leibold and Chase (2017), the relationship between taxonomic and functional  
103 homogenization allows us to infer which processes are acting: while taxonomic homogenization  
104 underlies dispersal limitation, functional homogenization is a signal of environmental filtering. In  
105 particular, anthropogenic environmental filters and spatial barriers are expected to differentially affect  
106 species within the original communities based on their functional traits. Therefore, exploring shifts  
107 in the functional profile of a community could shed light into mechanisms underlying community  
108 assemblages (Leibold and Chase, 2017).

109 We here analysed the community composition of diatoms in streams of the Eastern Italian Alps  
110 (Trento province), by collating a dataset of data collected during eight years of biomonitoring  
111 programs (2008-2016). In particular, we aimed at: i) investigating the differential role of  
112 environmental variables and spatial factors in driving the community composition in pristine and  
113 impacted sites within Alpine streams; and ii) unravelling whether taxonomic and/or functional  
114 homogenization occur in pristine and impacted sites and whether the lithological substrate may drive  
115 these processes. We hypothesized that: i) diatom composition in pristine sites would be mainly shaped  
116 by spatial factors, because the variation in chemical parameters within these oligotrophic ecosystems  
117 is expected to be low; conversely, chemical parameters would have a determinant role in impacted  
118 sites; ii) functional homogenization would occur in impacted sites as a consequence of environmental  
119 filters; and iii) lithological substrate would cause taxonomic differentiation in pristine sites, where  
120 species can track their ecological preferences without constraints of chemical variables (Rimet et al.,

2004; 2007; 2009; Tison et al., 2005; Grenier et al., 2006), but not in impacted sites, where chemical parameters are the main drivers of diatom community composition.

## **MATERIALS AND METHODS**

### **1.1. Study area**

The Trento province is located in the North-East of Italy and covers an area of nearly 6000 km<sup>2</sup>. Within this area, two hydroecoregions (hereafter HERs), i.e. homogeneous areas in terms of geology, altitude and climate (Wasson et al., 2002), were defined consistently with WFD requirements. Both HERs are Alpine and, within each HER, reference sites with potentially homogeneous, pristine environmental conditions and unaltered diatom communities were identified (Rimet et al., 2004). The first HER (hereafter calcareous HER), is the Southern Pre-Alps and Dolomites hydroecoregion, characterized by mountain streams flowing on calcareous substrate (massive and carbonated rocks). The second HER (hereafter siliceous HER) is the Inner Alps hydroecoregion, with high mountains and streams flowing on siliceous substrates (crystalline rocks). Diatom biomonitoring in streams has been carried out by the Environment Protection Agency of Trento province (APPA TN) since 2004. The monitoring network, including 166 sites in the entire Trento province, was designed to provide a coherent and comprehensive overview of ecological and chemical status within each river. In accordance with the monitoring program, every site is sampled twice in a year every six years (every three years for reference sites), considering biological, hydromorphological and physico-chemical quality elements, and pollutants discharged into the river basin or sub-basin.

In this study area, we analysed diatom communities collected from both HERs, in 17 reference and 16 impacted sites (Table 1), during the last eight years of biomonitoring programs (2008-2016). 11 reference sites were selected following the official national criteria (DM 56, 14/04/2009), i.e. sites characterized by null or slight human impacts, unaltered hydrological conditions, intact three-dimensional river connections, null or slight hydromorphological modifications, high water quality,

146 and biological communities showing taxonomical composition and densities corresponding to the  
147 unaltered conditions. Beside these official stations, 6 further sites with minor hydromorphological  
148 alterations were considered similar to reference sites because of their high water quality and were  
149 added to reference category for this study. Impacted sites are located in water bodies characterized  
150 by agricultural and urban pressures and they have high nutrient concentration and pesticide  
151 contamination (Fig. 1).

## 152 **1.2. Physical and chemical analyses**

153 Water samples for the physical and chemical analyses were collected along with diatom sampling. In  
154 both reference and impacted sites we analysed the following 15 parameters using standard methods  
155 (see Table 2): Biological Oxygen Demand (BOD<sub>5</sub>), carbonate hardness (HARD), Chemical Oxygen  
156 Demand (COD), Chlorides (Cl<sup>-</sup>), conductivity (COND), dissolved oxygen (DO), nitrates (N-NO<sub>3</sub>),  
157 orthophosphates (P-PO<sub>4</sub>), sulphates (SO<sub>4</sub>), temperature (TEMP), pH, total nitrogen (N<sub>tot</sub>), total  
158 phosphorous (P<sub>tot</sub>), total suspended sediments (TSS), turbidity (TURB).

## 159 **1.3. Diatom collection and treatment**

160 In total, we collected 64 diatom samples from 17 reference sites and 46 diatom samples from 16  
161 impacted sites (see Tab. 1 for details).

162 Samplings were performed between 2008 and 2016. Following the WFD requirements, each site  
163 belonging to surveillance monitoring was sampled at least twice during the 8 years monitoring  
164 program (mainly in spring and autumn). Some sites (Table A1), belonging to core network  
165 monitoring, were sampled in three different years (2 samples per year X 3 years = 6 samples).  
166 Epilithic diatoms were collected following the standard procedure (European Committee for  
167 Standardization, 2003). In each site, we chose 5 cobbles from the main flow and we collected  
168 periphyton by scraping their upper surface by means of a toothbrush. Samples were then preserved  
169 in ethanol and moved in laboratory for the treatment with H<sub>2</sub>O<sub>2</sub> (30%) and HCl (European Committee



for Standardization, 2003). Slides for the observation at the light microscope were mounted by means of Naphrax®. Diatom identification was based on several diatom floras and monographies, as well as recent taxonomic papers (Krammer and Lange-Bertalot, 1986-1991 a, b; Lange-Bertalot and Metzeltin, 1996; Krammer, 1997 a, b; 2002; 2003; Reichardt, 1999; Lange-Bertalot, 2001; Werum and Lange-Bertalot, 2004; Blanco et al., 2010; Hofmann et al. 2011; Bey and Ector, 2013; Falasco et al., 2013; Ector et al., 2015). We identified at least 400 valves in each sample. Diatom communities were analysed in terms of biodiversity, taxonomical composition, ecological guilds and growth forms (Rimet and Bouchez, 2012).

#### 1.4. Statistical analyses

All statistical analyses were performed with the software R 3.4.2 (R Development Core Team, 2017).

##### 1.4.1. Characterization of environmental features and diatom community

We performed the Mann-Whitney U-test to check for differences in environmental variables between references and impacted sites on the whole dataset and between the two HERs within the reference and impacted datasets. To highlight statistically significant species, representative of the two hydroecoregions for both reference and impacted sites, we performed an Indicator Species Analysis (Dufrêne and Legendre, 1997) with the function “multipatt” in the package *indicspecies* (Caceres and Legendre, 2009).

##### 1.4.2. Environmental parameters vs spatial processes

To test whether diatom communities in pristine sites are more shaped by spatial factors than in disturbed sites (Hypothesis 1), we investigated the role of environmental vs spatial parameters in both reference and impacted sites. To achieve this aim, we performed a partial-RDA, following the approach suggested in Peres-Neto et al. (2006) and De Bie et al. (2012). Using redundancy analysis (RDA), we built up two explanatory matrices: a chemical matrix [C], including all chemical parameters, and a spatial matrix [S] with both the coordinates of the sampling sites and the spatial variables extracted by Moran’s Eigenvector Maps analysis (MEM, see Dray et al., 2006). The MEM

analysis partition the spatial information into variables representing the potential autocorrelation between spatial points at different scales. With this procedure we generated a set of orthogonal, and thus non-collinear, spatial variables that are derived from geographical coordinates of the study sites, each of which corresponds to a specific spatial structure and scale. These variables can model coarse patterns in the community data and then progressively represent finer-scale patterns (Borcard et al., 2004). Both chemical and spatial variables were selected by means of a forward selection performed with the R package *packfor* (Dray et al., 2013) to obtain a parsimonious combination of variables, i.e. including only variables with a significant relationship with the community matrix. We separately tested the [C] and [S] matrices against the taxonomic matrix and we decomposed total community variation into pure components and their intersections. Significance was tested by means of a Monte Carlo test with 999 permutations.

#### 1.4.3. Taxonomic vs functional composition

To test whether anthropogenic pressure causes biotic homogenization in the taxonomic and/or the functional structure of diatom communities (Hypothesis 2), we analysed the taxonomic and functional response of diatom communities. First, we evaluated whether reference and impacted sites differed in their composition of diatom communities. Changes in taxonomic and functional composition (homogenization or differentiation) among the reference and impacted sites were analysed using the test of homogeneity for multivariate dispersion (Anderson et al., 2006) following the procedure proposed by Brice et al. (2017). This test represents a method to evaluate the homogenization within communities belonging to the same category and to compare these values among categories, e.g. among reference and impacted sites or among the calcareous and siliceous HERs. The test of homogeneity for multivariate dispersion was performed with the function “betadisper” in the package *vegan* (Oksanen et al., 2018). The taxonomic matrix, with the relative abundance of each recorded taxon in each sample, was converted into a site-by-site distance matrix using the Bray-Curtis distance, to which the test was applied. The distance of each site to its associated group multidimensional median was calculated and differences among such site distances were tested

221 by means of multivariate analogue of the Levene's test for homogeneity of variance with 9,999  
222 permutations to determine whether the dispersions between the two groups were different. We then  
223 repeated the test of homogeneity for multivariate dispersion on a functional matrix, containing trait  
224 abundances for each sampled site. To generate the functional matrix, we first created a species-by-  
225 trait matrix that was multiplied by the species-by-site matrix to obtain the site-by-trait matrix with  
226 the function "functcomp" in the package *FD* (Laliberté et al., 2014), in which each entry corresponds  
227 to the sum of the relative abundances of all the species present in a site that have a particular trait  
228 state. Functional traits considered for generating the functional matrix were life-forms, ecological  
229 guilds, size classes and biovolume (Rimet and Bouchez, 2012). We also tested shifts in taxonomic  
230 and functional composition between reference and impacted sites with a PERMANOVA (Anderson,  
231 2001) applied on distance matrices, using the function "adonis" in the package *vegan* (Oksanen et al.,  
232 2017). PERMANOVA is a non parametric multivariate analysis of variance that measures location  
233 differences in centroids of the different categories. Thus, we applied this test for investigating possible  
234 taxonomic and/or functional turnover among categories. Statistical significance was tested via 9999  
235 random permutations.

236 The same procedure was finally adopted to detect differences between the two HERs, which are  
237 proxies of lithological substrates, to test whether geology causes differentiation in diatom  
238 composition in pristine but not in impacted sites (Hypothesis 3).

#### 239 1.4.4. Analysis of functional traits

240 The set of functional metrics extrapolated from the functional matrix were subjected to a non-  
241 parametric Mann-Withtney U-test to test for differences between reference and impacted sites and  
242 between the calcareous and siliceous HERs within both reference and impacted sites.

243

## 244 2. RESULTS

### 245 2.1. Physical and chemical analyses

246 Observed values of physical and chemical parameters of reference sites showed low nutrient contents  
247 (Tab. 2) thus classifying them as oligotrophic in accordance with the Italian Water Legislation (D.  
248 Leg. 152/2006 and successive ones). BOD<sub>5</sub> concentrations never exceeded 2.5mg/L, being included  
249 in the range of the first water quality class of the national classification, like COD that on average  
250 never exceeded 3mg/L. DO was generally high, and pH ranged from 6.70 to 8.50, according to the  
251 riverbed lithology. Impacted sites generally showed lower quality, since, on average, N-NO<sub>3</sub> values  
252 were included in the range of the third water quality class (D. Leg. 152/2006 and successive ones),  
253 with mean values of 1.90 to 2.02 mg/L on calcareous and siliceous substrates respectively. Despite  
254 slightly higher than in reference sites, BOD<sub>5</sub> and COD levels corresponded to a moderate organic  
255 level (BOD<sub>5</sub> highest value = 5.1 mg/L; COD highest value = 13.8 mg/L) so as P<sub>tot</sub> whose maximum  
256 value reached 0.17 mg/L.

257 Reference and impacted sites showed significant differences in terms of all chemical parameters,  
258 except for temperature and TSS and all variables were higher in impacted than reference sites (Tab.  
259 2; Fig. B1). Among reference sites, results of the Mann-Whitney U-test showed that N-NO<sub>3</sub>, N<sub>tot</sub>, P-  
260 PO<sub>4</sub>, conductivity, hardness and pH were significantly higher in the calcareous than siliceous  
261 hydroecoregion, while opposite results were observed for BOD<sub>5</sub> and turbidity (Tab. 2; Fig. B2).  
262 Among impacted sites, we observed no significant differences between the two geological substrates  
263 (Tab. 2; Fig. B3).

## 264 **2.2. Diatom community composition**

265 In total, we analysed 64 diatom samples from reference sites and identified 120 taxa. In these samples,  
266 the most abundant and frequent species was *Achnantheidium minutissimum*, which was detected in all  
267 samples and represented the most abundant and frequent species, with a relative abundance on  
268 average of 35% per sample. *Achnantheidium pyrenaicum* (about 16% mean relative abundance),  
269 *Achnantheidium lineare* (9%). *Gomphonema pumilum* (8%), *Fragilaria arcus* (6%), *Encyonema*  
270 *minutum* (2%), *Fragilaria vaucheriae* (<2%), *Cocconeis lineata* (<2%), *Diatoma mesodon* (<2%)

271 were the most abundant taxa detected in at least 50% of the reference sites. In impacted sites, we  
272 analysed 46 diatom samples and identified 128 taxa. Again, *A. minutissimum* was the most abundant  
273 (mean relative abundance 12%) and frequent (detected in 98% of the samples) species. Beside *A.*  
274 *minutissimum*, 16 species were observed in at least 50% of the samples; among them, *Fistulifera*  
275 *saprophila* (about 11% of mean relative abundance), *Cocconeis euglypta* (6%), *Amphora pediculus*  
276 (6%), *Mayamaea permitis* (5%), *Nitzschia fonticola* (5%), *A. pyrenaicum* (3%), *Nitzschia*  
277 *inconspicua* (3%), *E. minutum* (3%), *Navicula tripunctata* (3%) and *Reimeria sinuata* (2%) were  
278 those exceeding 2% of relative abundance. According to the Indicator Species Analysis, several  
279 species are exclusive of the two HERs in pristine sites, with 10 species representative of the  
280 calcareous HER and 8 species representative of the siliceous HER (Tab. 3). Conversely, only few  
281 species are exclusive of the two HERs in impacted sites, with 2 species representative of the  
282 calcareous substrates and 4 species representative of the siliceous substrates (Tab. 3). *Fragilaria*  
283 *vaucheriae* is shared between reference and impacted sites as an indicator species for the siliceous  
284 HER, whereas *Gomphonema olivaceum* is representative of both siliceous reference sites and  
285 calcareous impacted sites.

### 286 **2.3. Environmental vs spatial components**

287 We first analysed whether diatom communities in pristine sites are more shaped by spatial factors than  
288 in disturbed sites (Hypothesis 1). The partial RDA performed on samples collected in reference sites  
289 (Fig. 2a) pointed out the minor role of chemical parameters, which together just explained 4% of the  
290 total explained variation. Only four chemical parameters (SO<sub>4</sub>, pH, N-NO<sub>3</sub> and DO) were included in  
291 the final environmental matrix after forward selection. On the other hand, the spatial matrix, which  
292 included seven PCNM vectors referring to both coarse and fine scale spatial autocorrelation after  
293 forward selection, and the spatially structured environmental parameters represented the key  
294 components in explaining diatom variability between the two HERs (8% and 12% respectively).

295 The partial-RDA performed on samples collected in impacted sites (Fig. 2b), highlighted again the  
296 dominant role of the spatial component, which alone explains 15% of the total variance. After forward  
297 selection, BOD, COD, Cl, HARD, N-NO<sub>3</sub> and SO<sub>4</sub> were included in the environmental matrix,  
298 whereas the spatial matrix included six PCNM vectors at large scale. Chemical parameters resulted  
299 important in shaping the communities of the impacted sites, explaining 10% of the total variance. The  
300 spatially structured environmental parameters explained 6% of the total variance.

#### 301 **2.4. Taxonomic vs functional composition**

302 Second, we analysed whether anthropogenic pressure causes taxonomic and/or functional  
303 homogenization of diatom communities (Hypothesis 2). Visual inspection of the PCoA ordination  
304 depicts a clear difference in both taxonomic and functional composition between references and  
305 impacted sites (Fig. 3). The results of the test of homogeneity for multivariate dispersion and  
306 PERMANOVA (Tab. 4) highlighted how reference and impacted sites show significant different  
307 taxonomic composition, with reference sites more homogeneous than impacted sites. However, even  
308 if the functional composition was significantly different between reference and impacted sites, we  
309 could not detect any homogenization effect. The analysis of the community functional profile (Fig.  
310 4) revealed significant higher abundances of *low* and *high profile* species in reference sites, whereas  
311 *motile* species were more abundant in impacted sites. Regarding life-forms, adnate species dominated  
312 in impacted sites, while pad-attached and stalked species showed significant higher abundances in  
313 reference sites.

314 Finally, we analysed whether lithological substrate causes biotic differentiation of diatom  
315 communities in pristine but not in impacted sites (Hypothesis 3). When considering reference sites,  
316 visual inspection of the PCoA performed on the taxonomic matrix clearly highlighted different  
317 species composition in siliceous and calcareous hydroecoregions (Fig. 5). This pattern was further  
318 confirmed by the results of the PERMANOVA (Table 4), which revealed significant differences in  
319 species composition between the two groups. Results of the test of homogeneity for multivariate

320 dispersion (Fig. 5; Tab. 4) highlighted that taxonomic and functional composition of diatom  
321 communities were homogeneous within the two groups, even if a nearly significant difference in  
322 terms of heterogeneity between diatom communities collected on siliceous (more heterogeneous) and  
323 calcareous (more homogeneous) hydroecoregions were observed in terms of ecological guilds. In  
324 particular, with the analysis of functional groups, we could detect how the *high profile* guild was  
325 significantly more abundant in the calcareous than siliceous area, while the *low profile* guild showed  
326 the opposite trend (Fig. 6). In terms of growth forms, the adnate, pad-attached and mucous-colonial  
327 taxa resulted significantly more abundant in the siliceous than calcareous hydroecoregion, whereas  
328 the opposite trend was observed for stalked-attached taxa (Fig. 6).

329 Concerning impacted sites, PCoA ordinations performed on the basis of both taxonomical and  
330 functional composition displayed an overlap between the two groups (Fig. 5). The test of  
331 homogeneity for multivariate dispersion underlined no significant differences between the two HERs,  
332 even if results of the PERMANOVA highlighted a slightly significant difference between the two  
333 groups in terms of taxonomic composition (Tab. 4). These results were furtherly corroborated by the  
334 Mann-Whitney U-test performed on total biovolume and relative abundance of ecological guilds and  
335 growth forms, which did not highlight differences between the two HERs (Fig. 6).

336

## 337 **DISCUSSION**

338 Traditionally, diatoms have been described as cosmopolitan organisms, whose communities are  
339 shaped by local environmental features (“everything is everywhere but environment selects”; Baas-  
340 Becking, 1934). For several years, this theory has been strengthened by the use of the *wide species*  
341 *concept* adopted in the European floras (Krammer and Large-Bertalot, 1986-1991), which led to the  
342 “force-fitting” of diatom taxa to “European” names and, as a consequence, to confusion concerning  
343 species distribution and diversity, which resulted strongly underestimated (Vanormelingen et al.,  
344 2008). However, in recent years, the role of spatial control, strictly associated to the dispersal

capacity, has been recognized as an important key factor for the determination of diatom community structure (Soininen, 2007), especially in Alpine streams (Bottin et al., 2014). We here investigated the differential role of environmental variables and spatial factors in driving the community composition in pristine and impacted sites within Alpine streams in NE-Italy. Our results confirmed that, in Alpine streams, differences in the composition of the communities are in part due to dispersal limitation. Previous studies in literature highlighted the importance of spatial variables, even at regional scale. For instance, in 2002, Potapova et Charles, observed that at national-scale (USA rivers), almost one-third of the variation in diatom species composition was explained by spatial factors, about 14% by spatially structured environmental parameters and more than 50% by local environmental features. Soininen (2007) highlighted that spatial component accounted 20-31% of the total variance explaining diatom community variations. However, the same author confirmed that at intermediate scale (i.e. 10-3000 km, corresponding to the scale of the present study), there is a joint control of both historical effects and contemporary ecological features. Dong et al. (2016) found that the spatial component played a significant role in shaping diatom communities (about 12%) even at small spatial extent (maximum distance between sites less than 30 km). In 2017, Marquardt and colleagues highlighted that the variation in compositional dissimilarity of some planktonic diatom communities collected in tropical reservoirs was better explained by geographic distance than local environmental features.

In accordance with Bottin et al. (2014), we found a stronger effect of the spatial component in impacted (15%) than in reference (8%) sites. This is in contrast with our initial hypothesis (Hypothesis 1) for which the importance of spatial component should be less evident in those sites affected by anthropogenic disturbance. In particular, we highlighted how environmental conditions in impacted sites likely cause biotic homogenization of diatom communities (Hypothesis 2), causing taxonomic but not functional homogenization. Therefore, the main functions in the community are maintained, while spatial processes likely eliminate some functional redundant species (Leibold and Chase, 2017). Spatial processes determining diatom community composition are controlled by



371 several factors, such as geographical barriers (i.e. spatial distance between sites), topographical and  
372 geomorphological features (i.e. high mountains), ecological traits of the species (i.e. their motility)  
373 but also by hydrological constraints, like artificial impoundments, which interrupt the longitudinal  
374 river connection (Dong et al., 2016). Even if the pressure of chemical parameters is evident in these  
375 streams, artificial alterations of hydrological connectivity likely play a major role in determining the  
376 structure of diatom community. Although the community spatial structure may be determined by  
377 multiple underlying mechanisms, like dispersal limitation, colonization stochasticity (i.e. priority  
378 effects) or individual spillover from upstream populations (i.e. mass effect), the analysis of the  
379 functional profile of diatom communities supports the role of dispersal limitation because the motile  
380 guild is dominant in impacted sites, which has been observed to be associated with both hydrological  
381 (Elias et al. 2015; Falasco et al. 2016) and morphological (Smucker and Vis, 2010; Bona et al. 2016)  
382 alterations. Our results therefore suggest that considering diatom response not only to trophic  
383 conditions but also to physical alterations is crucial in biomonitoring programs.

384 In reference sites, spatially structured environmental variables explain a great part of the total  
385 variance. This result suggests how chemical parameters strictly linked to peculiar areas (i.e. to HERs)  
386 are the main drivers of diatom community composition, further strengthening the role of geology. In  
387 fact, the different lithology characterizing the two hydroecoregions led to significant differences in  
388 some important geochemical variables, such as conductivity and hardness, which were higher on  
389 calcareous than on siliceous substrates (Rimet et al., 2004), with repercussions on diatom species  
390 composition. As already observed in previous studies, hydroecoregion environmental characteristics,  
391 like geology, resulted among the most important factors in shaping diatom reference communities of  
392 high altitude streams (Tison et al., 2005; Rimet et al., 2004; 2007). This is confirmed by our results,  
393 since in reference sites taxonomic composition of diatom communities significantly differed between  
394 the two HERs but this is not evident in impacted sites. This result confirms our third hypothesis,  
395 underlying that the influence of geology is much more important in reference than impacted sites,  
396 where the effect of geology is overridden by chemical parameters. The Indicator Species Analysis

397 confirmed such differences in terms of diatom communities in according with previous findings,  
 398 highlighting some species as significantly representative of reference conditions on calcareous  
 399 substrates, i.e. *Gomphonema pumilum* (Cantonati, 1998; Cantonati et al., 2012; Tison et al., 2005;  
 400 Rimet et al., 2004; 2007; Beltrami et al., 2012), *Gomphonema tergestinum* (Rimet et al., 2005; 2007;  
 401 Tison et al., 2005) and *Cymbella affinis* (Tison et al., 2003; 2005), and on siliceous substrates, i.e.  
 402 *Encyonema minutum* (Tison et al., 2005; Cantonati et al., 2012; Beltrami et al., 2012), *Encyonema*  
 403 *silesiacum* (Rimet et al., 2004; Bona et al., 2007; Beltrami et al., 2012), *Fragilaria vaucheriae* (Rimet  
 404 et al., 2004; Tison et al., 2005; Beltrami et al., 2012) and *Reimeria sinuata* (Rimet et al., 2004; Tison  
 405 et al., 2005; Beltrami et al., 2012).

406 We found contrasting results concerning *Gomphonema olivaceum*, which resulted indicator species  
 407 of siliceous substrates in our reference sites, while typical of the calcareous typologies in other studies  
 408 (see Rimet et al., 2004; Tison et al. 2005; Beltrami et al., 2012). However, in the present study *G.*  
 409 *olivaceum* also resulted as indicator species of calcareous substrates in the impacted typologies,  
 410 suggesting that probably geology is not the main parameter controlling the distribution and abundance  
 411 of this taxon. Even though *Achnantheidium lineare* did not result among the statistically significant  
 412 indicator species, we noticed it was mainly abundant on the calcareous hydroecoregion, confirming  
 413 previous findings in the study area (Beltrami et al., 2012).

414 Diatom communities in the two hydroecoregions within reference sites also differed in terms of  
 415 functional composition, confirming the significant functional shift obtained with the PERMANOVA.  
 416 In the siliceous sites, the *low profile* guild dominates the community, suggesting that the hydrological  
 417 regime differs between the two hydroecoregions. We can hypothesize that rivers flowing on the  
 418 siliceous substrates are possibly characterized by turbulent flows and unstable substrates, which did  
 419 not allow the development of a mature three-dimensional periphyton. Conversely, the calcareous  
 420 hydroecoregion presented more mature and stable communities, dominated by the *high profile* guild,  
 421 i.e. taxa which tolerate higher nutrient load, but sensitive to physical disturbance, suggesting more  
 422 stable flow conditions compared to siliceous streams (Rimet and Bouchez, 2012). Our hypothesis is

423 supported by Soininen (2007), who pointed out the role of flow velocity among the key environmental  
424 factors shaping diatom composition at local scale. In light of these results, the recent hydrological  
425 alteration that is affecting Alpine streams (Falasco et al. 2018; Piano et al. 2019) might represent a  
426 serious threat to diatom communities.

427 Conversely, differences in functional composition between HERs were less evident in impacted sites,  
428 confirming the absence of both functional homogenization and functional shift detected with the test  
429 of homogeneity for multivariate dispersion and PERMANOVA respectively.

430 Overall, our results are in accordance with the observations of Heino (2013) and highlight the  
431 possibility that water quality bioassessment based on diatoms should be considered more reliable at  
432 small scale. In fact, at drainage basin scale, the species sorting should be more effective than at larger  
433 scale, where the importance of the spatial component and dispersal limitation is typically high. As  
434 highlighted by Bottin et al. (2014), the effect of the spatial component on the structure of diatom  
435 assemblages can represent a background noise in the framework of the river quality assessment, and  
436 this should be taken into account especially in those countries, like Italy, covering a broad range of  
437 mountain areas. Given that physical barriers limiting dispersal in Alpine streams are mainly  
438 represented by orographic borders, the identification of reference sites within each drainage network  
439 could represent a solution to this constraint.

440 In addition, although we observed taxonomic homogenization in impacted sites, the turnover of  
441 species resulted to be more important than biotic homogenization in shaping diatom communities  
442 within Alpine streams. These results suggest that the variance between reference and impacted sites  
443 is comparable and that they are characterized by different species, thus strengthening the role of  
444 diatoms as bioindicators. In light of these patterns, the implementation of metrics that measure this  
445 component (e.g. Hillebrand et al., 2017) should be encouraged in biomonitoring programs.

## 446 **Acknowledgements**

447 This work is part of the research fellowship “From perennial to intermittent: structural and functional  
448 responses of benthic diatom communities in Alpine and Appenninic rivers” won by dr. Elisa Falasco.  
449 The authors declare no conflicts of interest.

450    **BIBLIOGRAPHY**

- 451    Anderson, M.J., 2001. A new method for non-parametric multivariate analysis of variance. *Austral.*  
452    *Ecol.* 26, 32-46.
- 453    Anderson, M. J., Ellingsen, K. E., McArdle, B. H., 2006. Multivariate dispersion as a measure of beta  
454    diversity. *Ecol. Lett.* 9(6), 683-693.
- 455    Baas Beeking, L.G.M., 1934. *Geobiologie of inleiding tot de milieukunde*. The Hague, the  
456    Netherlands: W.P. Van Stockum & Zoon (in Dutch).
- 457    Beltrami, M. E., Ciutti, F., Cappelletti, C., Lösch, B., Alber, R., Ector, L., 2012. Diatoms from Alto  
458    Adige/Südtirol (Northern Italy): characterization of assemblages and their application for biological  
459    quality assessment in the context of the Water Framework Directive. *Hydrobiologia* 695(1), 153-170.
- 460    Bey M. Y. & Ector L. (2013). *Atlas des diatomées des cours d'eau de la region Rhône-Alpes*. Tome  
461    1-6. Direction régionale de l'Environnement, de l'Aménagement et du Logement Rhône-Alpes.
- 462    Blanco S., Cejudo-Figueiras C., Álvarez-Blanco I., Bécares E., Hoffmann L., Ector L., 2010. *Atlas*  
463    *de las Diatomeas de la cuenca del Duero - Diatom Atlas of the Duero Basin*. 1a ed., Área de  
464    Publicaciones, Universidad de León, León, 382 pp.
- 465    Bona, F., Falasco, E., Fassina, S., Griselli, B., Badino, G., 2007. Characterization of diatom  
466    assemblages in mid-altitude streams of NW Italy. *Hydrobiologia* 583(1), 265-274.
- 467    Bona, F., Falasco, E., Fenoglio, S., Iorio, L., Badino, G., 2008. Response of macroinvertebrate and  
468    diatom communities to human-induced physical alteration in mountain streams. *River Res. App.*  
469    24(8), 1068-1081.
- 470    Bona, F., Doretto, A., Falasco, E., La Morgia, V., Piano, E., Ajassa, R., Fenoglio, S., 2016. Increased  
471    sediment loads in Alpine streams: an integrated field study. *River Res. App.* 32(6), 1316-1326.

472 Borcard, D., Legendre, P., Avois-Jacquet, C., Tuomisto, H., 2004. Dissecting the spatial structure of  
 473 ecological data at multiple scales. *Ecology* 85(7), 1826-1832.

474 Bottin, M., Soininen, J., Ferrol, M., Tison-Rosebery, J. 2014. Do spatial patterns of benthic diatom  
 475 assemblages vary across regions and years?. *Freshw. Sci.* 33(2), 402-416.

476 Brice, M. H., Pellerin, S., & Poulin, M. (2017). Does urbanization lead to taxonomic and functional  
 477 homogenization in riparian forests?. *Diversity and Distributions*, 23(7), 828-840.

478 Cáceres, M. D., Legendre, P., 2009. Associations between species and groups of sites: indices and  
 479 statistical inference. *Ecology* 90(12), 3566-3574.

480 Cantonati, M., 1998. Diatom communities of springs in the Southern Alps. *Diatom Res.* 13, 201-220.

481 Cantonati, M., Corradini, G., Juttner, I., Cox, J., 2001. Diatom assemblages in high mountain streams  
 482 of the Alps and the Himalaya. *Nova Hedwigia* 123, 37–62.

483 Cantonati, M., Gerecke, R., Bertuzzi, E., 2006. Springs of the Alps, sensitive ecosystems to  
 484 environmental change: From biodiversity assessments to long-term studies. In: Lami, A., Boggero,  
 485 A. (eds): *Developments of Hydrobiology, Ecology of high altitude aquatic systems in the Alps.*  
 486 *Hydrobiologia* 562, 59–96.

487 Cantonati, M., Decet, F., Corradini, F., Bertuzzi, E., 2007. The significance of chemical and physical  
 488 factors influencing the ecology of springs and a case study in the south-eastern Alps (Dolomiti  
 489 Bellunesi National Park). – In: Cantonati, M., Bertuzzi, E., Spitale, D. (eds): *The spring habitat: Biota*  
 490 *and sampling methods. Monografie del Museo Tridentino di Scienze Naturali*, 4, 47–78.

491 Cantonati, M., Spitale, D., 2009. The role of environmental variables in structuring epiphytic and  
 492 epilithic diatom assemblages in springs and streams of the Dolomiti Bellunesi National Park (south  
 493 eastern Alps). *Fundam. Appl. Limnol.* 174, 117-133.

494 Cantonati, M., Angeli, N., Bertuzzi, E., Spitale, D., Lange-Bertalot, H., 2012. Diatoms in springs of  
 495 the Alps: spring types, environmental determinants, and substratum. *Freshw. Sci.* 31, 499-524.

496 De Bie T., De Meester L., Brendonck L., Martens K., Goddeeris B., Ercken D., Hampel H., Denys  
 497 L., Vanhecke L., Van der Gucht K., Van Wichelen J., Vyverman W., Declerck S. A. J., 2012. Body  
 498 size and dispersal mode as key traits determining metacommunity structure of aquatic organisms.  
 499 *Ecol. Lett.* 15, 740–747.

500 Dong X.Y., Li B., He F.Z., Gu Y., Sun M.Q., Zhang H.M., Tan L., Xiao W., Liu S.R., Cai Q.H.,  
 501 2016. Flow directionality, mountain barriers and functional traits determine diatom metacommunity  
 502 structuring of high mountain streams. *Nature Sci. Rep.* 6, 24711.

503 Dray S., Legendre P., Peres-Neto P. R., 2006. Spatial modelling: a comprehensive framework for  
 504 principal coordinate analysis of neighbour matrices (PCNM). *Ecol. Modell.* 196, 483-493.

505 Dray S., Legendre P., Blanchet F. G., 2013. packfor: Forward selection with permutation. R package  
 506 version 0.0-8/r99.

507 Dufrêne, M., Legendre, P., 1997. Species assemblages and indicator species: the need for a flexible  
 508 asymmetrical approach. *Ecol. Monographs* 67(3), 345-366.

509 Ector L., Wetzel C. E., Novais M. H., Guillard D., 2015. Atlas des diatomées des rivières des Pays  
 510 de la Loire et de la Bretagne. DREAL Pays de la Loire, Nantes, 649pp.

511 Elias, C. L., Calapez, A. R., Almeida, S. F., Feio, M. J., 2015. From perennial to temporary streams:  
 512 an extreme drought as a driving force of freshwater communities' traits. *Mar. Freshw. Res.* 66(5),  
 513 469-480.

514 European Committee for Standardization, 2003. Water Quality Guidance Standard for the Routine  
 515 Sampling and Pretreatment of Benthic Diatoms from Rivers. European Standard EN 13946. European  
 516 Committee for Standardization, Brussels, 14 pp.

517 Falasco E., Piano E., Bona F., 2013. Guida al riconoscimento e all'ecologia delle principali diatomee  
518 fluviali dell'Italia nord occidentale. *Biol. Amb.* 27, 1-292.

519 Falasco, E., Ector, L., Ciaccio, E., Hoffmann, L., Bona, F., 2012. Alpine freshwater ecosystems in a  
520 protected area: a source of diatom diversity. *Hydrobiologia* 695(1), 233-251.

521 Falasco, E., Piano, E., Bona, F., 2016. Suggestions for diatom-based monitoring in intermittent  
522 streams. *Knowl. Manag. Aquat. Ecosyst.* (417), 38.

523 Falasco, E., Piano, E., Doretto, A., Fenoglio, S., Bona, F., 2018. Lentification in Alpine rivers:  
524 patterns of diatom assemblages and functional traits. *Aquat. Sci.* 80(4), 36.

525 Grenier, M., Campeau, S., Lavoie, I., Park, Y. S., Lek, S., 2006. Diatom reference communities in  
526 Quebec (Canada) streams based on Kohonen self-organizing maps and multivariate analyses. *Can. J.*  
527 *Fish. Aquat. Sci.* 63, 2087–2106.

528 Heino, J., 2013. The importance of metacommunity ecology for environmental assessment research  
529 in the freshwater realm. *Biol. Rev.* 88(1), 166-178.

530 Hillebrand, H., Blasius, B., Borer, E. T., Chase, J. M., Downing, J. A., Eriksson, B. K., Filstrup, C.  
531 T., Harpole, W. S., Hodapp, D., Larsen, S., Lewandowska, A. M., Seabloom, E. W., Van der Wall,  
532 D. B., Ryabov, A. B., 2018. Biodiversity change is uncoupled from species richness trends:  
533 consequences for conservation and monitoring. *J. Appl. Ecol.* 55(1), 169-184.

534 Hofmann, G., Werum, M., Lange-Bertalot, H., 2011. *Diatomeen im Süßwasser-Benthos von*  
535 *Mitteleuropa*. Koeltz Scientific Books, Königstein, 908 pp.

536 Krammer, K., 1997a. *Die cymbelloiden Diatomeen. Teil 1. Allgemeines und Encyonema Part.*  
537 *Bibliotheca Diatomologica* 36, 382 pp.



538 Krammer, K., 1997b. Die cymbelloiden Diatomeen. Teil 2. Encyonema part, Encyonopsis and  
539 Cymbelloopsis. Bibliotheca Diatomologica 37, 469 pp.

540 Krammer, K. 2002. Cymbella. In: Diatoms of Europe. 3., (Ed. H. Lange-Bertalot), A.R.G. Gantner  
541 Verlag KG, Rugell, 584pp.

542 Krammer, K., 2003. Cymbopleura, Delicata, Navicymbula, Gomphocymbelloopsis, Afrocymbella. In:  
543 Diatoms of Europe. 4. (Ed. H. Lange-Bertalot). ARG Gantner Verlag KG, Rugell, 530pp.

544 Krammer, K., Lange-Bertalot, H., 1986. Bacillariophyceae Teil: Naviculaceae. 1. In: Süßwasserflora  
545 von Mitteleuropa. 2. (Eds H. Ettl, J. Gerloff, H. Heynig & D. Mollenhauer). Fischer Verlag, Stuttgart,  
546 876 pp.

547 Krammer, K., Lange-Bertalot, H., 1988. Bacillariophyceae Teil: Bacillariaceae, Epithemiaceae,  
548 Surirellaceae. 2. In: Süßwasserflora von Mitteleuropa. 2. (Eds H. Ettl, J. Gerloff, H. Heynig & D.  
549 Mollenhauer). Fischer Verlag, Stuttgart, 610 pp.

550 Krammer, K., Lange-Bertalot, H., (1991a). Bacillariophyceae Teil: Centrales, Fragilariaceae,  
551 Eunotiaceae. 3. In: Süßwasserflora von Mitteleuropa. 2. (Eds H. Ettl, J. Gerloff, H. Heynig & D.  
552 Mollenhauer) Fischer Verlag, Stuttgart, 598 pp.

553 Krammer, K., Lange-Bertalot, H., 1991b. Bacillariophyceae Teil: Achnanthaceae. Kritische  
554 Ergänzungen zu Navicula (Lineolatae) und Gomphonema. 4. In: Süßwasserflora von Mitteleuropa.  
555 2. (Eds H. Ettl, J. Gerloff, H. Heynig & D. Mollenhauer). Fischer Verlag, Stuttgart, 437 pp.

556 Laliberté, E., Legendre, P., Shipley, B., 2014. FD: measuring functional diversity from multiple traits,  
557 and other tools for functional ecology. R package version 1.0-12.

558 Lange-Bertalot, H., 2001. *Navicula sensu stricto*, 10 Genera separated from *Navicula sensu lato*,  
559 *Frustulia*. In: *Diatoms of Europe*. 2. (Ed. H. Lange-Bertalot). A.R.G. Gantner Verlag K.G., Rugell,  
560 526 pp.

561 Lange-Bertalot H. and Metzeltin D., 1996. Indicators of Oligotrophy. In: Lange-Bertalot H (ed.)  
562 *Iconographia Diatomologica*. 2. Koeltz, Koenigstein, 390 p.

563 Leibold, M.A., Chase, J.M., 2018. *Metacommunity Ecology – Monographs in Population Biology*,  
564 Volume 59. Princeton University Press, 504 pp.

565 Marquardt, G.C., Blanco, S., Bicudo, C.E. de M., 2017. Distance decay as a descriptor of the diatom  
566 compositional variation in tropical reservoirs. *Mar. Freshw. Res.* 69, 105-113.

567 Oksanen, J., Guillaume Blanchet, F., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin,  
568 P.R., O'Hara, R.B., Simpson, G.L., Solymos, P., Henry, M., Stevens, H., Szoecs E., Wagner, H.,  
569 2018. *vegan: Community Ecology Package*. R package version 2.5-3.

570 Olden, J. D., Poff, N. L., Douglas, M. R., Douglas, M. E., Fausch, K. D., 2004. Ecological and  
571 evolutionary consequences of biotic homogenization. *TREE* 19(1), 18-24.

572 Peres-Neto, P. R., Legendre, P., Dray, S. Borcard, D., 2006. Variation partitioning of species data  
573 matrices: estimation and comparison of fractions. *Ecology* 87, 2614-2625.

574 Potapova, M.G., Charles, D.F.C., 2002. Benthic diatoms in USA rivers: distributions along spatial  
575 and environmental gradients. *J. Biogeogr.* 29,167–187.

576 R Core Team, 2018. *R: A language and environment for statistical computing*. R Foundation for  
577 Statistical Computing, Vienna, Austria. <http://www.R-project.org/>.

578 Reichardt, E., 1999. Zur Revision der Gattung Gomphonema. Die Arten um *G. affine/insigne*, *G.*  
579 *angustum/micropus*, *G. acuminatum* sowie gomphonemoide Diatomeen aus dem Oberoligozän in

580 Böhmen. In: *Iconographia Diatomologica*. 8, (Ed. H. Lange-Bertalot) A.R.G. Gantner Verlag K.G.,  
581 Rugell,

582 Rimet, F., 2009. Benthic diatom assemblages and their correspondence with ecoregional  
583 classifications: case study of rivers in north-eastern France. *Hydrobiologia* 636, 137–151.

584 Rimet, F., Bouchez, A., 2012. Life-forms, cell-sizes and ecological guilds of diatoms in European  
585 rivers. *Knowl. Manag. Aquat. Ecosyst.* 406, 01-12.

586 Rimet, F., Ector, L., Cauchie, H. M., Hoffmann, L., 2004. Regional distribution of diatom  
587 assemblages in the headwater streams of Luxembourg. *Hydrobiologia* 520, 105-117.

588 Rimet, F., Cauchie, H. M., Tudesque, L. & Ector, L., 2005. Use of artificial intelligence (MIR-max)  
589 and chemical index to define type diatom assemblages in Rhhe basin and Mediterranean region. In:  
590 Lek, S., Scardi, M., Verdonchot, P.F.M., Descy, J.P., Park, Y.S. (eds.) *Modelling community*  
591 *structure in aquatic ecosystems*. Springer Verlag, pp. 288-303.

592 Rimet, F., Gomà, J., Cambra, J., Bertuzzi, E., Cantonati, M., Cappelletti, C., Ciutti, F., Cordonier, A.,  
593 Coste, M., Delmas, F., Tison, J., Tudesque, L., Vidal, H., Ector, L., 2007. Benthic diatoms in western  
594 European streams with altitudes above 800 m: characterisation of the main assemblages and  
595 correspondence with ecoregions, *Diat. Res.* 22:1, 147-188.

596 Smucker, N. J., Vis, M. L., 2010. Using diatoms to assess human impacts on streams benefits from  
597 multiple-habitat sampling. *Hydrobiologia* 654(1), 93-109.

598 Soininen, J., 2007. Environmental and spatial control of freshwater diatoms – a review. *Diat. Res.*  
599 22(2), 473-490.

600 Southwood, T. R., 1977. Habitat, the templet for ecological strategies?. *J. Anim. Ecol.* 46(2), 337-  
601 365.

602 Southwood, T. R., 1988. Tactics, strategies and templets. *Oikos* 3-18.

603 Tison, J., Coste, M., Delmas, F., Giraudel, J. L., Lek, S., 2003. Typologie des assemblages  
604 diatomiques du bassin Adour-Garonne (France) par l'utilisation combinee d'un reseau artificiel de  
605 neurones (algorithme de Kohonen) et de l'indice de structuration. *Bul. Soc. Nat. Ouest Fr.* 2, 16-229.

606 Tison, J., Park, Y. S., Coste, M., Wasson, J. G., Ector, L., Rimet, F., Delmas, F., 2005. Typology of  
607 diatom communities and the influence of hydro-ecoregions: a study on the French hydrosystem scale.  
608 *Wat. Res.* 39(14), 3177-3188.

609 Townsend, C. R., Hildrew, A. G., 1994. Species traits in relation to a habitat templet for river systems.  
610 *Freshw. Biol.* 31(3), 265-275.

611 Vanormelingen, P., Verleyen, E., Vyverman, W., 2008. The diversity and distribution of diatoms:  
612 from cosmopolitanism to narrow endemism. *Biodiv. Cons.* 17(2), 393-405.

613 Verleyen, E., Vyverman, W., Sterken, M., Hodgson, D. A., De Wever, A., Juggins, S., Van der Vijver,  
614 B., Jones, V.J., Vanormelingen, P., Roberts, D., Flower, R., Kilroy, C., Souffreau, C., Sabbe, K.,  
615 2009. The importance of dispersal related and local factors in shaping the taxonomic structure of  
616 diatom metacommunities. *Oikos* 118(8), 1239-1249.

617 Vilmi, A., Karjalainen, S.M., Hellsten, S., Heino, J., 2016. Bioassessment in a metacommunity  
618 context: are diatom communities structured solely by species sorting? *Ecol. Ind.* 62, 86-94.

619 Ward, J. V., 1994. Ecology of Alpine streams. *Freshw. Biol.* 32(2), 277-294.

620 Wasson, J. G., Chandesris, A., Pella, H., Blanc, L., 2002. Typology and reference conditions for  
621 surface water bodies in France: the hydro-ecoregion approach. *TemaNord* 566, 37-41.

622 Werum, M., Lange-Bertalot, H., 2004. Diatoms in springs from Central Europe and elsewhere under  
623 the influence of hydrogeology and anthropogenic impacts. In: *Iconographia Diatomologica* 13. (Ed.  
624 H. Lange-Bertalot) Koeltz, Koenigstein: 417 pp.

625

626 **Tables**

627 Table 1. Characterization of sampling sites (CAL = calcareous HER; SIL = siliceous HER).

SITES	HER	origin	distance from the source/morphology	influence of the upstream basin	n samples	n total
REFERENCE SITES	CAL	intermittent	meandering	null	2	64
		spring	< 10 km		2	
		pluvial/nival	< 5 km		12	
			5-25 km		4	
			25-75 km	slight	4	
	SIL	glacial	< 10 km	not detectable	6	
		pluvial/nival	< 5 km		6	
			5-25 km		28	
IMPACTED SITES	CAL	pluvial/nival	< 5 km	null	4	46
			5-25 km	strong	16	
			25-75 km		8	
			slight	slight	2	
		subterranean	< 10 km	null	2	
	SIL	pluvial/nival	1.24	not detectable	6	
			0.9		8	

628

629 Table 2. average (mean), standard deviation (sd), minimum (min) and maximum (max) values of chemical parameters observed in the calcareous  
630 HER (CAL) and in the siliceous HER (SIL) in both reference and impacted sites (N-NO<sub>3</sub> = nitrates; N\_TOT = total nitrogen; P-PO<sub>4</sub> =  
631 orthophosphates; P\_TOT = total phosphorous; SO<sub>4</sub> = sulphates; Cl<sup>-</sup> = chlorides; BOD<sub>5</sub> = biochemical oxygen demand; COD = chemical oxygen  
632 demand; HARD = hardness; COND = conductivity; DO = dissolved oxygen; TEMP = temperature; TSS = total suspended solids; TURB =  
633 turbidity).

Sites	HER	value	N-NO <sub>3</sub> (mg/l)	N_TOT (mg/l)	P-PO <sub>4</sub> (mg/l)	P_TOT (mg/l)	SO <sub>4</sub> (mg/l)	Cl <sup>-</sup> (mg/l)	BOD <sub>5</sub> (mg/l)	COD (mg/l)	HARD (mg/l)	COND (μS/cm)	DO (mg/l)	pH	TEMP (°C)	TSS (mg/l)	TURB (NTU)
REFERENCE SITES	CAL	mean	0.700	0.680	0.008	0.010	5.60	0.670	0.521	2.09	132	220	10.9	8.24	8.00	5.00	1.12
		sd	0.200	0.260	0.008	0.012	4.30	0.470	0.102	1.10	36.7	47	0.720	0.190	2.80	0.00	0.954
		min	0.400	0.250	0.005	0.005	1.40	0.200	0.500	2.50	17.0	128	9.40	7.70	4.30	5.00	0.300
		max	1.12	1.28	0.040	0.040	16.4	2.00	1.00	6.20	168	277	12.7	8.40	15.0	5.00	4.20
	SIL	mean	0.330	0.390	0.005	0.012	17.8	0.940	0.80	2.70	51.0	100	10.7	7.60	7.50	12.0	7.12
		sd	0.160	0.230	0.001	0.016	17.6	1.12	0.500	0.800	37.7	66	0.940	0.390	2.80	24.0	17.7
		min	0.100	0.250	0.005	0.005	0.800	0.100	0.500	2.50	5.00	14	9.00	6.70	2.80	5.00	0.200
		max	0.900	1.10	0.010	0.060	58.0	6.20	2.30	5.90	139	250	12.6	8.50	14.0	129	98.0
IMPACTED SITES	CAL	mean	2.02	2.29	0.059	0.045	13.4	7.25	1.30	4.30	190	339	11.4	8.39	9.10	6.00	4.24
		sd	1.21	1.38	0.038	0.035	9.90	4.19	1.00	2.80	63.3	99.0	1.25	0.220	4.00	4.00	4.25
		min	0.500	0.600	0.010	0.005	2.60	0.900	0.500	2.50	42.0	92.0	9.40	7.80	1.00	5.00	0.600
		max	7.50	8.40	0.070	0.160	43.0	19.4	5.10	13.8	291	531	15.2	8.80	17.3	25.0	22.3
	SIL	mean	1.90	2.13	0.056	0.070	42.7	10.5	1.10	5.00	133	267	11.4	8.07	7.80	5.00	3.30
		sd	1.26	1.24	0.031	0.037	70.2	3.19	0.700	2.40	97.3	150	0.960	0.160	3.40	0.00	2.88
		min	0.80	0.900	0.020	0.020	5.30	6.80	0.500	2.50	74.0	157	10.1	7.90	3.00	5.00	1.00
		max	4.00	4.20	0.110	0.130	238	16.5	2.90	9.20	398	975	12.9	8.30	13.6	5.00	11.2

634

635 Table 3. Species resulting indicators of calcareous (CAL) and siliceous (SIL) HER in reference and  
636 impacted sites from the Indicator Species Analysis

SITES	CAL			SIL		
	species	IndVal	P	species	IndVal	P
REFERENCE SITES	<i>Gomphonema pumilum</i>	0.817	0.001	<i>Encyonema minutum</i>	0.844	0.001
	<i>Denticula tenuis</i>	0.678	0.001	<i>Fragilaria vaucheriae</i>	0.831	0.001
	<i>Diatoma erhenbergii</i>	0.640	0.006	<i>Encyonema silesiacum</i>	0.716	0.002
	<i>Cymbella excisa</i>	0.589	0.001	<i>Gomphonema olivaceum</i>	0.699	0.002
	<i>Gomphonema tergestinum</i>	0.520	0.020	<i>Reimeria sinuata</i>	0.686	0.038
	<i>Nitzschia fonticola</i>	0.519	0.039	<i>Psammothidium bioretii</i>	0.539	0.020
	<i>Gomphonema elegantissimum</i>	0.497	0.006	<i>Gomphonema pumilum</i> var. <i>rigidum</i>	0.474	0.026
	<i>Cocconeis placentula</i> var. <i>placentula</i>	0.462	0.018	<i>Achananthidium atomoides</i>	0.447	0.034
	<i>Cymbella affinis</i>	0.424	0.040			
	<i>Cymbella excisiformis</i>	0.405	0.023			
IMPACTED SITES	<i>Gomphonema olivaceum</i>	0.712	0.010	<i>Nitzschia inconspicua</i>	0.814	0.002
	<i>Encyonema ventricosum</i>	0.707	0.004	<i>Fragilaria vaucheriae</i>	0.679	0.046
				<i>Rhoicosphenia abbreviata</i>	0.600	0.020
				<i>Diatoma mesodon</i>	0.513	0.029

637

638




639 Table 4. Results of the test for differences in species composition between reference and impacted  
640 sites and between the calcareous (CAL) and the siliceous (SIL) HER within both reference and  
641 impacted sites as inferred from the test of homogeneity for multivariate dispersion and  
642 PERMANOVA analysis. Significant values are highlighted in bold.

	<b>Taxonomic composition</b>	<b>Functional composition</b>
<b>Reference vs impacted sites</b>		
Homogeneity test	$F_{1,108} = 16.2$ <b><math>P &lt; 0.001</math></b>	$F_{1,108} = 0.140$ $P = 0.709$
PERMANOVA	$F_{1,108} = 24.1$ <b><math>P = 0.001</math></b>	$F_{1,108} = 49.9$ <b><math>P = 0.001</math></b>
<b>CAL vs SIL in reference sites</b>		
Homogeneity test	$F_{1,62} = 0.315$ $P = 0.577$	$F_{1,62} = 3.93$ $P = 0.052$
PERMANOVA	$F_{1,62} = 7.32$ <b><math>P = 0.001</math></b>	$F_{1,62} = 4.81$ <b><math>P = 0.002</math></b>
<b>CAL vs SIL in impacted sites</b>		
Homogeneity test	$F_{1,44} = 0.194$ $P = 0.662$	$F_{1,44} = 0.598$ $P = 0.444$
PERMANOVA	$F_{1,44} = 1.79$ <b><math>P = 0.047</math></b>	$F_{1,44} = 1.92$ $P = 0.110$

643


644



646

647 Figure 1. Map of the sampling sites (purple circles = reference sites; brown circles = impacted sites)  
 648 in the study area (light blue = calcareous HER; orange = siliceous HER).


649



650

651 Figure 2. Results from variation partitioning (partial redundancy analysis). The relative contributions  
652 (% of explanation) of chemical and spatial variables, as well as the shared components explaining  
653 variation in diatom communities in reference (a) and impacted (b) sites.

654




655

656 Figure 3. PCoA ordinations of samples based on taxonomic (left panel) and functional (right panel)


657 composition of diatom communities in reference (purple) and impacted (brown) sites.

658




659

660 Figure 4. Boxplots depicting the relative abundances of ecological guilds and life-forms in reference  
 661 (purple) and impacted (brown) sites. Results of the Mann-Whitney U-test are reported below each  
 662 graph. Asterisks refer to significant differences between the two categories (\* < 0.05; \*\* < 0.01; \*\*\*  
 663 < 0.001).



664

665 Figure 5. PCoA ordinations of samples based on taxonomic (left panel) and functional (right panel)  
 666 composition of diatom communities in calcareous (light blue) and siliceous (orange) sites performed  
 667 on reference (upper panel) and impacted (lower panel) sites.



668

669 Figure 6. Boxplots depicting the relative abundances of ecological guilds and life-forms in calcareous  
 670 (light blue) and siliceous (orange) substrates in reference (upper panel) and impacted (lower panel)  
 671 sites. Asterisks refer to significant differences between the two categories (\* < 0.05; \*\* < 0.01; \*\*\*  
 672 < 0.001).